

Table C.1: Heat Capacities of Gases in the Ideal-Gas State[†]

Constants in equation $C_P^{ig}/R = A + BT + CT^2 + DT^{-2}$ T (kelvins) from 298 to $T_{\max}$

Chemical species		$T_{\max}$	C_P^{ig}/R	A	$10^3 B$	$10^6 C$	$10^{-5} D$
Paraffins:							
Methane	CH ₄	1500	4.217	1.702	9.081	-2.164	
Ethane	C ₂ H ₆	1500	6.369	1.131	19.225	-5.561	
Propane	C ₃ H ₈	1500	9.011	1.213	28.785	-8.824	
<i>n</i> -Butane	C ₄ H ₁₀	1500	11.928	1.935	36.915	-11.402	
<i>iso</i> -Butane	C ₄ H ₁₀	1500	11.901	1.677	37.853	-11.945	
<i>n</i> -Pentane	C ₅ H ₁₂	1500	14.731	2.464	45.351	-14.111	
<i>n</i> -Hexane	C ₆ H ₁₄	1500	17.550	3.025	53.722	-16.791	
<i>n</i> -Heptane	C ₇ H ₁₆	1500	20.361	3.570	62.127	-19.486	
<i>n</i> -Octane	C ₈ H ₁₈	1500	23.174	4.108	70.567	-22.208	
1-Alkenes:							
Ethylene	C ₂ H ₄	1500	5.325	1.424	14.394	-4.392	
Propylene	C ₃ H ₆	1500	7.792	1.637	22.706	-6.915	
1-Butene	C ₄ H ₈	1500	10.520	1.967	31.630	-9.873	
1-Pentene	C ₅ H ₁₀	1500	13.437	2.691	39.753	-12.447	
1-Hexene	C ₆ H ₁₂	1500	16.240	3.220	48.189	-15.157	
1-Heptene	C ₇ H ₁₄	1500	19.053	3.768	56.588	-17.847	
1-Octene	C ₈ H ₁₆	1500	21.868	4.324	64.960	-20.521	
Miscellaneous organics:							
Acetaldehyde	C ₂ H ₄ O	1000	6.506	1.693	17.978	-6.158	
Acetylene	C ₂ H ₂	1500	5.253	6.132	1.952		-1.299
Benzene	C ₆ H ₆	1500	10.259	-0.206	39.064	-13.301	
1,3-Butadiene	C ₄ H ₆	1500	10.720	2.734	26.786	-8.882	
Cyclohexane	C ₆ H ₁₂	1500	13.121	-3.876	63.249	-20.928	
Ethanol	C ₂ H ₆ O	1500	8.948	3.518	20.001	-6.002	
Ethylbenzene	C ₈ H ₁₀	1500	15.993	1.124	55.380	-18.476	
Ethylene oxide	C ₂ H ₄ O	1000	5.784	-0.385	23.463	-9.296	
Formaldehyde	CH ₂ O	1500	4.191	2.264	7.022	-1.877	
Methanol	CH ₄ O	1500	5.547	2.211	12.216	-3.450	
Styrene	C ₈ H ₈	1500	15.534	2.050	50.192	-16.662	
Toluene	C ₇ H ₈	1500	12.922	0.290	47.052	-15.716	
Miscellaneous inorganics:							
Air		2000	3.509	3.355	0.575		-0.016
Ammonia	NH ₃	1800	4.269	3.578	3.020		-0.186
Bromine	Br ₂	3000	4.337	4.493	0.056		-0.154
Carbon monoxide	CO	2500	3.507	3.376	0.557		-0.031
Carbon dioxide	CO ₂	2000	4.467	5.457	1.045		-1.157
Carbon disulfide	CS ₂	1800	5.532	6.311	0.805		-0.906
Chlorine	Cl ₂	3000	4.082	4.442	0.089		-0.344
Hydrogen	H ₂	3000	3.468	3.249	0.422		0.083
Hydrogen sulfide	H ₂ S	2300	4.114	3.931	1.490		-0.232
Hydrogen chloride	HCl	2000	3.512	3.156	0.623		0.151
Hydrogen cyanide	HCN	2500	4.326	4.736	1.359		-0.725
Nitrogen	N ₂	2000	3.502	3.280	0.593		0.040
Nitrous oxide	N ₂ O	2000	4.646	5.328	1.214		-0.928
Nitric oxide	NO	2000	3.590	3.387	0.629		0.014
Nitrogen dioxide	NO ₂	2000	4.447	4.982	1.195		-0.792
Dinitrogen tetroxide	N ₂ O ₄	2000	9.198	11.660	2.257		-2.787
Oxygen	O ₂	2000	3.535	3.639	0.506		-0.227
Sulfur dioxide	SO ₂	2000	4.796	5.699	0.801		-1.015
Sulfur trioxide	SO ₃	2000	6.094	8.060	1.056		-2.028
Water	H ₂ O	2000	4.038	3.470	1.450		0.121

[†] Selected from H. M. Spencer, *Ind. Eng. Chem.*, vol. 40, pp. 2152-2154, 1948; K. K. Kelley, *U.S. Bur. Mines Bull.* 584, 1960; L. B. Pankratz, *U.S. Bur. Mines Bull.* 672, 1982.

Table C.4: Standard Enthalpies and Gibbs Energies of Formation
at 298.15 K[†]

Joules per mole of the substance formed

Chemical species		State (Note 2)	$\Delta H_{f,298}^{\circ}$ (Note 1)	$\Delta G_{f,298}^{\circ}$ (Note 1)
Paraffins:				
Methane	CH ₄	(g)	-74,520	-50,460
Ethane	C ₂ H ₆	(g)	-83,820	-31,855
Propane	C ₃ H ₈	(g)	-104,680	-24,290
<i>n</i> -Butane	C ₄ H ₁₀	(g)	-125,790	-16,570
<i>n</i> -Pentane	C ₅ H ₁₂	(g)	-146,760	-8,650
<i>n</i> -Hexane	C ₆ H ₁₄	(g)	-166,920	150
<i>n</i> -Heptane	C ₇ H ₁₆	(g)	-187,780	8,260
<i>n</i> -Octane	C ₈ H ₁₈	(g)	-208,750	16,260
1-Alkenes:				
Ethylene	C ₂ H ₄	(g)	52,510	68,460
Propylene	C ₃ H ₆	(g)	19,710	62,205
1-Butene	C ₄ H ₈	(g)	-540	70,340
1-Pentene	C ₅ H ₁₀	(g)	-21,280	78,410
1-Hexene	C ₆ H ₁₂	(g)	-41,950	86,830
1-Heptene	C ₇ H ₁₄	(g)	-62,760	
Miscellaneous organics:				
Acetaldehyde	C ₂ H ₄ O	(g)	-166,190	-128,860
Acetic acid	C ₂ H ₄ O ₂	(l)	-484,500	-389,900
Acetylene	C ₂ H ₂	(g)	227,480	209,970
Benzene	C ₆ H ₆	(g)	82,930	129,665
Benzene	C ₆ H ₆	(l)	49,080	124,520
1,3-Butadiene	C ₄ H ₆	(g)	109,240	149,795
Cyclohexane	C ₆ H ₁₂	(g)	-123,140	31,920
Cyclohexane	C ₆ H ₁₂	(l)	-156,230	26,850
1,2-Ethanediol	C ₂ H ₆ O ₂	(l)	-454,800	-323,080
Ethanol	C ₂ H ₆ O	(g)	-235,100	-168,490
Ethanol	C ₂ H ₆ O	(l)	-277,690	-174,780
Ethylbenzene	C ₈ H ₁₀	(g)	29,920	130,890
Ethylene oxide	C ₂ H ₄ O	(g)	-52,630	-13,010
Formaldehyde	CH ₂ O	(g)	-108,570	-102,530
Methanol	CH ₄ O	(g)	-200,660	-161,960
Methanol	CH ₄ O	(l)	-238,660	-166,270
Methylcyclohexane	C ₇ H ₁₄	(g)	-154,770	27,480
Methylcyclohexane	C ₇ H ₁₄	(l)	-190,160	20,560
Styrene	C ₈ H ₈	(g)	147,360	213,900
Toluene	C ₇ H ₈	(g)	50,170	122,050
Toluene	C ₇ H ₈	(l)	12,180	113,630

Table C.4 (Continued)

Chemical species		State (Note 2)	ΔH_{f298}° (Note 1)	ΔG_{f298}° (Note 1)
Miscellaneous inorganics:				
Ammonia	NH ₃	(g)	-46,110	-16,450
Ammonia	NH ₃	(aq)		-26,500
Calcium carbide	CaC ₂	(s)	-59,800	-64,900
Calcium carbonate	CaCO ₃	(s)	-1,206,920	-1,128,790
Calcium chloride	CaCl ₂	(s)	-795,800	-748,100
Calcium chloride	CaCl ₂	(aq)		-8,101,900
Calcium chloride	CaCl ₂ ·6H ₂ O	(s)	-2,607,900	
Calcium hydroxide	Ca(OH) ₂	(s)	-986,090	-898,490
Calcium hydroxide	Ca(OH) ₂	(aq)		-868,070
Calcium oxide	CaO	(s)	-635,090	-604,030
Carbon dioxide	CO ₂	(g)	-393,509	-394,359
Carbon monoxide	CO	(g)	-110,525	-137,169
Hydrochloric acid	HCl	(g)	-92,307	-95,299
Hydrogen cyanide	HCN	(g)	135,100	124,700
Hydrogen sulfide	H ₂ S	(g)	-20,630	-33,560
Iron oxide	FeO	(s)	-272,000	
Iron oxide (hematite)	Fe ₂ O ₃	(s)	-824,200	-742,200
Iron oxide (magnetite)	Fe ₃ O ₄	(s)	-1,118,400	-1,015,400
Iron sulfide (pyrite)	FeS ₂	(s)	-178,200	-166,900
Lithium chloride	LiCl	(s)	-408,610	
Lithium chloride	LiCl·H ₂ O	(s)	-712,580	
Lithium chloride	LiCl·2H ₂ O	(s)	-1,012,650	
Lithium chloride	LiCl·3H ₂ O	(s)	-1,311,300	
Nitric acid	HNO ₃	(l)	-174,100	-80,710
Nitric acid	HNO ₃	(aq)		-111,250
Nitrogen oxides	NO	(g)	90,250	86,550
	NO ₂	(g)	33,180	51,310
	N ₂ O	(g)	82,050	104,200
	N ₂ O ₄	(g)	9,160	97,540
Sodium carbonate	Na ₂ CO ₃	(s)	-1,130,680	-1,044,440
Sodium carbonate	Na ₂ CO ₃ ·10H ₂ O	(s)	-4,081,320	
Sodium chloride	NaCl	(s)	-411,153	-384,138
Sodium chloride	NaCl	(aq)		-393,133
Sodium hydroxide	NaOH	(s)	-425,609	-379,494
Sodium hydroxide	NaOH	(aq)		-419,150
Sulfur dioxide	SO ₂	(g)	-296,830	-300,194
Sulfur trioxide	SO ₃	(g)	-395,720	-371,060
Sulfur trioxide	SO ₃	(l)	-441,040	
Sulfuric acid	H ₂ SO ₄	(l)	-813,989	-690,003
Sulfuric acid	H ₂ SO ₄	(aq)		-744,530
Water	H ₂ O	(g)	-241,818	-228,572
Water	H ₂ O	(l)	-285,830	-237,129

* From *TRC Thermodynamic Tables—Hydrocarbons*, Thermodynamics Research Center, Texas A & M Univ. System, College Station, TX: "The NBS Tables of Chemical Thermodynamic Properties," *J. Phys. and Chem. Reference Data*, vol. 11, supp. 2, 1982.

Notes

1. The standard property changes of formation ΔH_{f298}° and ΔG_{f298}° are the changes occurring when 1 mol of the listed compound is formed from its elements with each substance in its standard state at 298.15 K (25°C).
2. Standard states: (a) Gases (g): pure ideal gas at 1 bar and 25°C. (b) Liquids (l) and solids (s): pure substance at 1 bar and 25°C. (c) Solutes in aqueous solution (aq): Hypothetical ideal 1-molal solution of solute in water at 1 bar and 25°C.