

EBP Implementation Scale

Below are 18 questions about evidence-based practice (EBP). Some healthcare providers do some of these things more often than other healthcare providers. There is no certain frequency in which you should be performing these tasks. Please answer each question by circling the number that best describes how often each item has applied to you in the past 8 weeks.

In the past 8 weeks, I have:

	0 times	1-3 times	4-5 times	6-7 times	8 or more times
1. Used evidence to change my clinical practice...	0	1	2	3	4
2. Critically appraised evidence from a research study...	0	1	2	3	4
3. Generated a PICO question about my clinical practice...	0	1	2	3	4
4. Informally discussed evidence from a research study with a colleague...	0	1	2	3	4
5. Collected data on a patient problem...	0	1	2	3	4
6. Shared evidence from a study or studies in the form of a report or presentation to more than 2 colleagues...	0	1	2	3	4
7. Evaluated the outcomes of a practice change...	0	1	2	3	4
8. Shared an EBP guideline with a colleague...	0	1	2	3	4
9. Shared evidence from a research study with a patient/family member...	0	1	2	3	4
10. Shared evidence from a research study with a multi-disciplinary team member...	0	1	2	3	4
11. Read and critically appraised a clinical research study...	0	1	2	3	4
12. Accessed the Cochrane Database of Systematic Reviews...	0	1	2	3	4
13. Accessed the National Guidelines Clearinghouse...	0	1	2	3	4
14. Used an EBP search or systematic review to change clinical practice where I work...	0	1	2	3	4
15. Evaluated a care initiative by collecting patient outcome data...	0	1	2	3	4
16. Shared the outcome data collected with colleagues...	0	1	2	3	4
17. Changed practice based on patient outcome data...	0	1	2	3	4
18. Promoted the use of EBP to my colleagues...	0	1	2	3	4

Copyright, Melnyk & Fineout-Overholt, 2003. Please DO NOT USE this instrument without permission from the authors. For further information about use, please contact arccle2003@gmail.com. Validity of this scale has been established and Cronbach alphas have been $\geq .85$ across various samples. For detailed information on the validity and reliability of this instrument, please see: Melnyk, B.M., Fineout-Overholt, E., & Mays, M. (2008). The evidence-based practice beliefs and implementation scales: Psychometric properties of two new instruments. *Worldviews on Evidence-Based Nursing*, 5(4), 208-216.